

Authors' contributions

Huizhong Wang: The study concept and design; literature search; data collection; critical review of the literature; writing of the manuscript or critical review of important intellectual; final approval of the final version of the manuscript.

Jingru Sun: The study concept and design; critical review of the literature; writing of the manuscript or critical review of important intellectual; final approval of the final version of the manuscript.

Conflicts of interest

None declared.

References

1. Raposo I, Torres T. Palmoplantar psoriasis and palmoplantar pustulosis: current treatment and future prospects. *Am J Clin Dermatol.* 2016;17:349–58.
2. Misiak-Galazka M, Zozula J, Rudnicka L. Palmoplantar Pustulosis: recent advances in etiopathogenesis and emerging treatments. *Am J Clin Dermatol.* 2020;21:355–70.
3. Maverakis E, Marzano AV, Le ST, Callen JP, Bruggen MC, Guenova E, et al. Pyoderma gangrenosum. *Nat Rev Dis Primers.* 2020;6:81.
4. McPhie ML, Kirchhof MG. Pyoderma gangrenosum treated with secukinumab: a case report. *SAGE Open Med Case Rep.* 2020;8, 2050313X20940430.
5. Coe J, Kudva S, Shams K. Matching the dose to the disease: successful treatment of recalcitrant pyoderma gangrenosum using high dose secukinumab. *Dermatol Ther.* 2022;35:e15669.
6. Orita A, Hoshina D, Hirosaki K. Pyoderma gangrenosum caused by secukinumab successfully treated with risankizumab: a case report and literature review. *Clin Exp Dermatol.* 2022;47:1372–4.
7. Jin K, Matsuzaki Y, Akasaka E, Nakano H, Sawamura D. Pyoderma gangrenosum triggered by switching from adalimumab to secukinumab. *J Dermatol.* 2019;46:e108–9.
8. Wollina U, Schönlebe J, Furr C. Pyoderma gangrenosum induced by secukinumab—a late paradoxical drug reaction. *Dermatol Ther.* 2020;33:e13161.
9. Bodis G, Toth V, Schwarting A. Role of human leukocyte antigens (HLA) in autoimmune diseases. *Rheumatol Ther.* 2018;5:5–20.
10. Yamamoto T. Similarity and difference between palmoplantar pustulosis and pustular psoriasis. *J Dermatol.* 2021;48:750–60.

Huizhong Wang  ^{a,b,c,d}, Jingru Sun  ^{a,b,c,d,*}

^a Department of Dermatology and Venereology, Peking University First Hospital, Beijing, China

^b Beijing Key Laboratory of Molecular Diagnosis on Dermatoses, Beijing, China

^c National Clinical Research Center for Skin and Immune Diseases, Beijing, China

^d NMPA Key Laboratory for Quality Control and Evaluation of Cosmetics, Beijing, China

* Corresponding author.

E-mail: sjr12315@126.com (J. Sun).

Received 25 April 2023; accepted 18 June 2023

Available online 2 September 2024

<https://doi.org/10.1016/j.abd.2023.06.012>

0365-0596/ © 2024 Sociedade Brasileira de Dermatologia.

Published by Elsevier España, S.L.U. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Squamous cell carcinoma of the nail unit after repeated UV nail lamp exposure. A call for action?*



Dear Editor,

An otherwise healthy 26-year-old female consulted Dermatology Service due to 1 year of changes in the color of her nail, and detachment of the nail plate on the fourth left finger. The patient referred had been continuously exposed to UV nail lamps twice a month for two years. She did not use sunscreen or any other form of protection while using the device. She did not use tanning beds either. She had a negative mycologic test performed and received topical mycological treatments with no response.

On examination, proximal leukonychia, distal yellow-brown chromonychia, and onychomadesis were observed (Fig. 1).

A biopsy of the nail matrix was performed, and the histopathology showed hyperparakeratosis, acanthosis, and intraepidermal proliferation of atypical keratinocytes, absence of maturation, and abundant mitoses. A squamous cell carcinoma in situ with partial resection (Fig. 2). Due to



Figure 1 Proximal leukonychia, distal yellow-brown chromonychia, and onychomadesis on the fourth left finger.

* Study conducted at the Hospital Italiano de Buenos Aires, Buenos Aires, Argentina.




Figure 2 Histopathology. Hyperparakeratosis and intraepidermal proliferation of atypical keratinocytes with abundant mitoses (Hematoxylin & eosin, 400×).



Figure 3 Postoperative result. A scar on the nail bed is observed.

the high-risk tumor location, Mohs micrographic surgery was indicated (Fig. 3).

Squamous cell carcinoma is a malignant tumor of keratinizing cells in the epidermis and its appendages. Well-defined risk factors exist for its development, with the main one being exposure to Ultraviolet Radiation (UVR).¹ Within the UVR spectrum, type A and its association with squamous cell carcinoma is well-known after prolonged exposure.²

Currently, gel or acrylic manicure is a very common practice in the population that requires UVA with fluorescent or LED lamps that have an emission spectrum of 375 to 425 nanometers.³ Currently, some case reports suggest a link between the use of UVA nail lamps and the development of squamous cell carcinomas and actinic keratoses, either on fingers or hands dorsum.^{4–7} The literature and case reports published so far conclude that the risks are potential and are limited to giving recommendations on their use. In a recently published study on Nature Communications, the molecular effect of the radiation emitted by UV nail lamps was experimentally evaluated in mammalian and human cells, demonstrating that it is cytotoxic, genotoxic, and mutagenic, predisposing to an increased risk of carcinomas.²

Some authorities such as the Food and Drug Administration and the World Health Organization mention the risk of carcinomas due to exposure to ultraviolet radiation, either

from the sun or artificial sources such as tanning beds. Still, they do not mention lamps used for manicures.^{8,9}

With this case presentation, while we can't assert a declared relationship between the use of UV lamps and the development of squamous cell carcinoma in our patient, it was highly suggestive due to being a young woman without other risk factors, with the presence of a single presentation tumor in an area directly exposed to nail-curing lamps.

In light of permanent manicures and acrylic nails becoming increasingly popular, we consider it important to analyze the issue due to the potential impact this practice could have, especially on young people unaware of the possible risks of this habit.

Financial support

None declared.

Authors' contributions

Tatiana Ordoñez: The study concept and design; Writing of the manuscript.

Marina Ruf: The study concept and design; Intellectual participation in the propaedeutic and/or therapeutic conduct of the studied cases.

Valeria Angles: Data collection, or analysis and interpretation of data.

Gabriel Brau: Writing of the manuscript or critical review of important intellectual content.

Damián Ferrario: Data collection.

Luis Mazzuccolo: Final approval of the final version of the manuscript.

Conflicts of interest

None declared.

References

- Zhivagui M, Hoda A, Valenzuela N, Yeh YY, Dai J, He Y, et al. DNA damage and somatic mutations in mammalian cells after irradiation with a nail polish dryer. *Nat Commun.* 2023;14:276.
- Shihab N, Lim HW. Potential cutaneous carcinogenic risk of exposure to UV nail lamp: a review. *Photodermatol Photoimmunol Photomed.* 2018;34:362–5.
- MacFarlane DF, Alonso CA. Occurrence of nonmelanoma skin cancers on the hands after UV nail light exposure. *Arch Dermatol.* 2009;145:447–9.
- Ratycz MC, Lender JA, Gottwald LD. Multiple dorsal hand actinic keratoses and squamous cell carcinomas: a unique presentation following extensive UV nail lamp use. *Case Rep Dermatol.* 2019;11:286–91.
- O'Sullivan NA, Tait CP. Tanning bed and nail lamp use and the risk of cutaneous malignancy: a review of the literature. *Australas J Dermatol.* 2014;55:99–106.
- Freeman C, Hull C, Sontheimer R, Curtis J. Squamous cell carcinoma of the dorsal hands and feet after repeated exposure to ultraviolet nail lamps. *Dermatol Online J.* 2020;26, 13030/qt1rd1k82v.
- Williams EA, Kuschner SH. Ultraviolet nail lamps and squamous cell carcinoma: cause and effect, or not? – A case report. *Open Journal of Orthopedics.* 2021;11:335–9.

8. Office of the Commissioner. How to Safely Use Nail Care Products [Internet]. U.S. Food and Drug Administration. FDA; [cited 2023 Feb 14]. Available from: <https://www.fda.gov/consumers/consumer-updates/how-safely-use-nail-care-products>.
9. World Health Organization [Internet]. Ultraviolet radiation. [cited 2023 Feb 14]. Available from: <https://www.who.int/es/news-room/fact-sheets/detail/ultraviolet-radiation>

Tatiana Ordoñez *, Marina Ruf , Valeria Angles , Gabriel Brau , Damián Ferrario , Luis Mazzuccolo 

Dermatology Department, Hospital Italiano de Buenos Aires, Buenos Aires, Argentina

* Corresponding author.

E-mail: tatiana.ordonez@hospitalitaliano.org.ar
(T. Ordoñez).

Received 31 May 2023; accepted 14 July 2023
Available online 6 August 2024

<https://doi.org/10.1016/j.abd.2023.07.020>

0365-0596/ © 2024 Sociedade Brasileira de Dermatologia.

Published by Elsevier España, S.L.U. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Vascularized parallel-ridge pattern: dermoscopic sign in acral melanoma with anatomopathological correlation[☆]



Dear Editor,

Acral lentiginous melanoma is a rare subtype of melanoma that affects the palms, soles and nails. Its early diagnosis is challenging, mainly due to a higher proportion of amelanotic melanomas and a wide variety of clinical presentations.¹ Patients often present with advanced disease at the time of diagnosis and therefore have a worse prognosis when compared to other melanoma subtypes.² Dermoscopy of the lesions shows specific patterns, being of great value for an early and accurate diagnosis. This is a case report of a recently described dermoscopic sign.

A 61-year-old woman reported the appearance of a brownish lesion on the right plantar region three years before, with progressive growth and ulceration. She had the

lesion excised at other hospital, but the material was not sent for anatomopathological study. She was then referred to ours hospital.

On clinical examination, she had a surgical scar in good aspect, measuring approximately 3 cm, on the right plantar region. Contact dermoscopy with polarized light of the skin adjacent to the scar showed erythema and dotted vessels filling the ridges and sparing the furrows (Fig. 1) – a recently described dermoscopic pattern, called “vascularized parallel-ridge pattern”³

The clinical history and the identification of the vascularized parallel-ridge pattern led to a high suspicion of melanoma. Therefore, a wider resection with a 2-cm margin was performed. Histopathology showed residual melanoma in the deep dermis (Figs. 2 and 3) and surgical margins free of neoplasia. In correspondence with the dermoscopic changes there were grouped proliferated and dilated capillary vessels close to the eccrine ducts (Fig. 4).

Acral lentiginous melanomas can show characteristic dermoscopic patterns of great diagnostic value. For pigmented lesions, three patterns should be highlighted: the parallel-ridge pattern with a sensitivity of 86% and specificity of 99%;




Figure 1 Patient's right plantar region, in the first evaluation at the hospital. On the left, the clinical photo shows the surgical scar from the previous excision at another medical service. On the right, contact dermoscopy with polarized light of the skin adjacent to the scar highlights the “vascularized parallel-ridge pattern” in the area circled in black.

[☆] Study conducted at the Department of Dermatology, Escola Paulista de Medicina, Universidade Federal de São Paulo, São Paulo, SP, Brazil.